

GENERALIZED M^* -SIMPLE GROUPS

SEBAHATTIN IKIKARDES AND RECEP SAHIN

ABSTRACT. Let X be a compact bordered Klein surface of algebraic genus $p \geq 2$, and let $G = \Gamma^*/\Lambda$ be a group of automorphisms of X where Γ^* is an NEC group and Λ is a bordered surface group. If the order of G is $4q/(q-2)(p-1)$, for $q \geq 3$ a prime number, then the signature of Γ^* is $(0; +; [-]; \{(2, 2, 2, q)\})$. These groups of automorphisms are called generalized M^* -groups. In this paper, we define generalized M^* -simple groups and give some examples of them. Also, we classify solvable generalized M^* -simple groups.

1. Introduction. A compact bordered Klein surface X of algebraic genus $p \geq 2$ admits at most $12(p-1)$ automorphisms [10]. Groups isomorphic to the automorphism group of such a compact bordered Klein surface with this maximal number of automorphisms are called M^* -groups. These groups were first introduced in [11], and have been studied in several papers ([3–5, 9]). Also, the survey article in [5] contains a nice survey of known results about M^* -groups.

An important result about M^* -groups is that they must have a certain partial presentation. This is established by considering an M^* -group as an epimorphic image of a quadrilateral group $\Gamma^*[2, 2, 2, 3]$. A quadrilateral group Γ^* is a non-Euclidean crystallographic (NEC) group with signature

$$(0; +; [-]; \{(2, 2, 2, 3)\}).$$

Also Γ^* is isomorphic to the abstract group with the presentation

$$\langle c_0, c_1, c_2, c_3 \mid c_i^2 = (c_0c_1)^2 = (c_1c_2)^2 = (c_2c_3)^2 = (c_3c_0)^3 = I \rangle.$$

For some bordered surface group Λ the group $G = \Gamma^*/\Lambda$ satisfies $|G| = 12(p-1)$ and there is a bordered smooth epimorphism $\theta : \Gamma^* \rightarrow G$

2010 AMS *Mathematics subject classification.* Primary 30F50, 20H10.
Keywords and phrases. Bordered Klein surface, automorphism, NEC group, M^* -group, generalized M^* -group, generalized M^* -simple group.
Received by the editors on September 14, 2009, and in revised form on June 16, 2010.

which maps $c_0 \rightarrow r_1$, $c_1 \rightarrow I$, $c_2 \rightarrow r_2$ and $c_3 \rightarrow r_3$. It is clear that $\text{Ker}(\theta) = \Lambda$. Thus, $r_1 r_2$ and $r_1 r_3$ have orders 2 and 3, respectively, and each group G admits the following partial presentation:

$$\langle r_1, r_2, r_3 \mid r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^2 = (r_1 r_3)^3 = \dots = I \rangle.$$

In [12, Proposition 2, page 223], May extended this in the following proposition to an extended quadrilateral group $\Gamma^*[2, 2, 2, n]$, $n \geq 3$, integer.

Proposition 1.1. *Let G be a finite group, and let $\Gamma^* = \Gamma^*[2, 2, 2, n]$ be an extended quadrilateral group. If there is a homomorphism $\phi : \Gamma^* \rightarrow G$ onto G such that $K = \ker \phi$ is a bordered surface group, then G is generated by three distinct nontrivial elements r_1 , r_2 and r_3 satisfying the relations*

$$(1.1) \quad r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^2 = (r_1 r_3)^n = I.$$

It is clear that the order of G is $[4n/(n-2)](p-1)$, where $p \geq 2$ is an integer, and the signature of Γ^* is $(0; +; [-; \{(2, 2, 2, n)\}])$.

In [12], for $n \geq 3$ a prime number, Sahin et al. referred to these finite groups, which were obtained by May in [12, Proposition 2, page 223], as *generalized M^* -groups*. A generalized M^* -group associated to $n \geq 3$ a prime number, is a finite group G generated by three distinct nontrivial elements r_1 , r_2 and r_3 which satisfy the relations (1.1). Notice that, if $n = 3$, then generalized M^* -groups are M^* -groups.

On the other hand, in [16, 17] the extended Hecke group $\overline{H}(\lambda_q)$ has been defined by adding the reflection $R(z) = 1/\bar{z}$ to the generators of the Hecke group $H(\lambda_q)$, where $q \geq 3$ is an integer, and it has been extensively studied (for examples, see [1, 8] and [6, page 70]). The extended Hecke group $\overline{H}(\lambda_q)$ has the presentation

$$\overline{H}(\lambda_q) = \langle R_1, R_2, R_3 \mid R_1^2 = R_2^2 = R_3^2 = (R_1 R_2)^2 = (R_1 R_3)^q = I \rangle,$$

where $R_1 = 1/\bar{z}$, $R_2 = -\bar{z}$ and $R_3 = -\bar{z} - \lambda_q$, $\lambda_q = 2 \cos(\pi/q)$ and $q \geq 3$ is an integer. The signature of the extended Hecke group $\overline{H}(\lambda_q)$ is $(0; +; [-; \{(2, q, \infty)\}])$. Since the extended Hecke group $\overline{H}(\lambda_q)$ contain

a reflection, it is an NEC group. Thus, the quotient space $\mathcal{U}/\overline{H}(\lambda_q)$ is a Klein surface and $\mathcal{U}/H(\lambda_q)$ is the canonical double cover of $\mathcal{U}/\overline{H}(\lambda_q)$ where \mathcal{U} is the upper half-plane. If a bordered surface group Γ is a normal subgroup of finite index in $\overline{H}(\lambda_q)$, then $\overline{H}(\lambda_q)/\Gamma$ is a group of automorphisms of the compact bordered Klein surface $X = \mathcal{U}/\overline{H}(\lambda_q)$. Also, the automorphism groups G of order $[4q/(q-2)](p-1)$ which act on compact bordered Klein surfaces X of genus $p \geq 2$, are finite quotient groups of the extended Hecke groups $\overline{H}(\lambda_q)$, where $q \geq 3$ is an integer. For example, the groups of orders $|G| = 12(p-1)$, $|G| = 8(p-1)$, $|G| = (20/3)(p-1)$, respectively, are the finite quotient groups of the extended Hecke groups $\overline{H}(\lambda_3)$ (the extended modular group $\text{PGL}(2, \mathbf{Z})$), $\overline{H}(\lambda_4)$ or $\overline{H}(\lambda_5)$ [20]. Here the orders of these groups are the highest three among the automorphism groups of the compact Klein surfaces of genus $p \geq 2$ (see [12, page 221, Proposition 1]).

In [11], May showed that there is a relationship which says a finite group of order at least 12 is an M^* -group if and only if it is a finite homomorphic image of the extended modular group.

It is easy to see that there is a relationship between extended quadrilateral groups $\Gamma^*[2, 2, 2, n]$, $n \geq 3$ integer and extended Hecke groups $\overline{H}(\lambda_q)$, $q \geq 3$ integer. Of course, there is a relationship between extended Hecke groups $\overline{H}(\lambda_q)$, $q \geq 3$ prime, and generalized M^* -groups (see the following diagram).

$$\begin{aligned} \overline{H}(\lambda_3) = \text{PGL}(2, \mathbf{Z}) &\longleftrightarrow M^*\text{-groups} \\ \overline{H}(\lambda_q) &\longleftrightarrow \text{generalized } M^*\text{-groups.} \end{aligned}$$

In fact, many results can be obtained by using these relations. As a consequence, in [18], Sahin et al. show that a finite group of order at least $4q$ is a generalized M^* -group if and only if it is the homomorphic image of the extended Hecke group $\overline{H}(\lambda_q)$. By using known results about normal subgroups of the extended Hecke groups $\overline{H}(\lambda_p)$ given in [19], they obtained an infinite family of generalized M^* -groups. Also, using known results about commutator subgroups of $\overline{H}(\lambda_p)$, the authors obtained that if G is a generalized M^* -group, then $|G : G'|$ divides 4 and $|G' : G''|$ divides q^2 . Finally, they proved that, if $q \geq 3$ is a prime number and G is a generalized M^* -group associated to q , then G is supersolvable if and only if $|G| = 4 \cdot q^r$ for some positive integer r .


In this paper, we define generalized M^* -simple groups and give some examples of them. Also, we classify solvable generalized M^* -simple groups.

2. Generalized M^* -simple groups. In this section, we want to define generalized M^* -simple groups in a manner analogous to M^* -simple groups, O^* -simple groups and $LO1$ -simple groups for the automorphism groups of Klein and Riemann surfaces given by May et al. in [9, 13, 14], respectively. To do this, we need the following theorem first.

Theorem 2.1. *Let $q \geq 3$ be a prime number, and let G be a generalized M^* -group associated to an extended quadrilateral group $\Gamma^*[2, 2, 2, q]$, with genus action on the bordered Klein surface X of genus $g \geq 2$. Let N be a normal subgroup of G of index $r > 2q$. Set $G' = G/N$, $X' = X/N$, let $\phi : X \rightarrow X'$ be the quotient map, and let g' be the genus of X' . Then:*

- (1) $g' \geq 2$;
- (2) G' is a generalized M^* -group;
- (3) ϕ is a full covering.

Proof. Firstly, we prove that $g' \geq 2$. The quotient space X/G is the disc D , and the quotient map $\pi : X \rightarrow D$ is ramified at four points, all of ∂D , with ramification indices $2, 2, 2, q$, where $q \geq 3$ is a prime number. From the induced action of $G' = G/N$ on $X' = X/N$, we have the following diagram of quotient maps.



Applying the Riemann-Hurwitz formula to the mapping ν yields

$$2g' - 2 = r \left[-2 + \sum \left(1 - \frac{1}{e_i} \right) \right],$$

where the e_i 's are chosen from $\{2, 2, 2, q\}$. Then it is easy to see that there are no solutions for $g' = 0$ or 1 , $r > 2q$. Thus, $g' \geq 2$.

Let G have generators r_1, r_2 and r_3 satisfying

$$r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^2 = (r_1 r_3)^q = I,$$

and let $\mu : G \rightarrow G' = G/N$ be a natural quotient map since the index $r = [G : N] > 2q$. It is easy to see that $r_1, r_2, r_3, r_1 r_2$ and $r_1 r_3$ are not in N . Then G' is generated by $\mu(r_1), \mu(r_2)$ and $\mu(r_3)$, and therefore G' is a generalized M^* -group.

Now, if we apply the Riemann-Hurwitz formula to the mapping ϕ , we find $2g' - 2 = |G'|(q - 2)/8q$ since $|G'| = |G/N|$ is the index in the NEC-group Γ^* of the obvious smooth epimorphism from Γ^* to G' . Then $|G'| = [4q/(q - 2)](g' - 1)$. Therefore, ϕ is unramified. \square

This theorem leads to the following notion.

Definition 2.1. A generalized M^* -group is called a generalized M^* -simple group if it has no non-trivial normal subgroups of index greater than $2q$, or equivalently, if it has no proper generalized M^* -quotient groups.

If a generalized M^* -group G has a quotient group of order $2q$ or less (the possibilities being the trivial group, $C_2, C_2 \times C_2, C_q$ and $C_2 \times C_q$), then these quotient groups are not generalized M^* -groups.

On the other hand, a simple generalized M^* -group is a generalized M^* -simple group. Thus, if G is a simple generalized M^* -group, then G acts only on non-orientable surfaces since otherwise the orientation-preserving maps would be subgroup of index 2 in G [10, page 206].

Now we give some examples related to generalized M^* -simple groups.

Example 2.1. Let q be an odd prime. The group $C_2 \times D_q \cong D_{2q}$ is the smallest generalized M^* -simple group. Let $D_{2q} = \langle A, B \mid A^2 = B^2 = (AB)^{2q} = I \rangle$. If we choose $r_1 = (AB)^s A, r_2 = B(AB)^s$ and $r_3 = B(AB)^t$, where $s = (q - 1)/2$ and $t = (q - 3)/2$, then D_{2q} has the following relations:

$$r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^2 = (r_1 r_3)^q = (r_2 r_3)^{2q} = (r_1 r_2 r_3)^2 = I.$$

Thus it is easy to check that the group D_{2q} is a generalized M^* -simple group.

Example 2.2. Let q be an odd prime. Let $G^{q,n,r}$ be the group with generators A, B and C and defining relations:

$$A^q = B^n = C^r = (AB)^2 = (BC)^2 = (CA)^2 = (ABC)^2 = I.$$

If we set $r_1 = BC$, $r_2 = CA$ and $r_3 = BCA$, then we obtain the presentation:

$$r_1^2 = r_2^2 = r_3^2 = (r_1 r_2)^2 = (r_1 r_3)^q = (r_2 r_3)^n = (r_1 r_2 r_3)^r = I.$$

Thus, G is a quotient of $\Gamma^*[2, 2, 2, q]$ by a bordered surface group if and only if G is a quotient of the group $G^{q,n,r}$ for some n and r . If $q \geq 3$ prime and the group is finite, then we obtain a generalized M^* -group with index n . Some values of n and r which make the group finite are given in [7]. It is clear that, if (q', n', r') is any permutation of (q, n, r) , then $G^{q',n',r'}$ is isomorphic to $G^{q,n,r}$. For $q = 7$, the groups $G^{7,9,3}$ and $G^{7,12,3}$ are generalized M^* -simple groups.

Example 2.3. Let q be an odd prime. The group $\text{PSL}(2, q)$ when $q \equiv 1 \pmod{4}$ and the group $\text{PGL}(2, q)$ when $q \equiv 3 \pmod{4}$ have the following presentation:

$$A^3 = B^{n(q)} = C^q = (AB)^2 = (BC)^2 = (CA)^2 = (ABC)^2 = I,$$

where $n(q)$ is the ordinal of the first Fibonacci number that is divisible by q ([3, page 54] and [9, page 277]). Thus, for the above values of q , the groups $\text{PSL}(2, q)$ and $\text{PGL}(2, q)$ are generalized M^* -simple groups.

It is hard to classify all generalized M^* -simple groups, or even simple generalized M^* -groups. In fact, classifying which simple groups are generalized M^* -groups is equivalent to classifying which simple groups are quotients of the extended Hecke groups $\overline{H}(\lambda_q)$. But to study these groups is very difficult. Here we can only classify solvable generalized M^* -simple groups.

Theorem 2.2. Let $q \geq 3$ be a prime number, and let G be a solvable generalized M^* -simple group associated to q .

- i) If $q = 3$, then $G \cong C_2 \times S_3$ or $G \cong S_4$;
- ii) If $q > 3$ then $G \cong C_2 \times D_q$.

Proof. i) Please see the proof in [9, Theorem 15, page 278].

ii) Let $q > 3$ be a prime number. If we check the groups of order $4qs$ where $s = 1, 2, \dots, q$, then we obtain that the two smallest solvable generalized M^* -groups are $C_2 \times D_q$ and $D_q \times D_q$ of orders $4q$ and $4q^2$, respectively. But the group $D_q \times D_q$ is not a generalized M^* -simple group. Then we assume that G is a solvable generalized M^* -group with $o(G) > 4q^2$. If we show that G has a generalized M^* -quotient group, then the proof will be complete.

We know that G is solvable. Then $G \neq G'$. Therefore, using Corollary 11 in [18, page 1215], we find $[G : G''] \geq 2q$. Also, since $o(G) > 4q^2$, we have $G'' \neq 1$. Now we consider two cases. If $[G : G'''] > 2q$, then using Theorem 2.1, we obtain that the quotient group G/G'' is a generalized M^* -group. Now we consider the case $[G : G'''] = 2q$, that is, $G/G'' \cong D_q$. If G'' were a minimal normal subgroup of G , then, from [15, Theorem 5.24, page 105], G'' would be an elementary Abelian 2-group. Also, G'' has no elements with order larger than $2q$. Thus, G'' is a quotient group of a group $G^{q,m,n}$, where $m \leq n \leq 2q$. But if we check the table in [7, pages 138–139] for each $q > 3$, then we see that there is no such generalized M^* -group G such that $[G : G'''] = 2q$ and $o(G) > 4q^2$. Hence, G'' contains a minimal normal subgroup N of G with $[G : N] > [G : G'''] = 2q$. Therefore, the quotient group G/N is a generalized M^* -group. \square

REFERENCES

1. F. Ates and A.S. Cevik, *Knit products of some groups and their applications*, Rend. Sem. Mat. Univ. Padova **121** (2009), 1–11.
2. E. Bujalance, F.J. Cirre and P. Turbek, *Groups acting on bordered Klein surfaces with maximal symmetry*, in *Proceedings of Groups St. Andrews 2001 in Oxford*, Vol. 1, London Math. Soc. Lect. Note Ser. **304**, Cambridge Univ. Press, Cambridge, 2003.
3. ———, *Subgroups of M^* -groups*, Quart. J. Math. **54** (2003), 49–60.
4. ———, *Automorphism criteria for M^* -groups*, Proc. Edinb. Math. Soc. **47** (2004), 339–351.
5. E. Bujalance, J.J. Etayo, J.M. Gamboa and G. Gromadzki, *Automorphism groups of compact bordered Klein surfaces. A combinatorial approach*, Lect. Notes Math. **1439**, Springer Verlag, 1990.
6. I.N. Cangül and D. Singerman, *Normal subgroups of Hecke groups and regular maps*, Math. Proc. Camb. Phil. Soc. **123** (1998), 59–74.
7. H.S.M. Coxeter and W.O.J. Moser, *Generators and relations for discrete groups*, 3rd ed., Ergeb. Math. Grenz. **14**, Springer-Verlag, Berlin, 1972.

8. M.L.A.N. de las Peñas, R.P. Felix and M.C.B. Decena, *Enumeration of index 3 and 4 subgroups of hyperbolic triangle symmetry groups*, *Z. Krist.* **223** (2008), 543–551.
9. N. Greenleaf and C.L. May, *Bordered Klein surfaces with maximal symmetry*, *Trans. Amer. Math. Soc.* **274** (1982), 265–283.
10. C.L. May, *Automorphisms of compact Klein surfaces with boundary*, *Pacific J. Math.* **59** (1975), 199–210.
11. ———, *Large automorphism groups of compact Klein surfaces with boundary*, *Glasgow Math. J.* **18** (1977), 1–10.
12. ———, *The groups of real genus 4*, *Mich. Math. J.* **39** (1992), 219–228.
13. ———, *The real genus of groups of odd order*, *Rocky Mountain J. Math.* **37** (2007), 1251–1269.
14. C.L. May and J. Zimmerman, *The symmetric genus of groups of odd order*, *Houston J. Math.* **34** (2008), 319–338.
15. J.J. Rotman, *An introduction to the theory of groups*, Fourth edition, *Grad. Texts Math.* **148**, Springer-Verlag, New York, 1995.
16. R. Sahin and O. Bizim, *Some subgroups of the extended Hecke groups $\overline{H}(\lambda_q)$* , *Acta Math. Sci.* **23** (2003), 497–502.
17. R. Sahin, O. Bizim and I.N. Cangül, *Commutator subgroups of the extended Hecke groups $\overline{H}(\lambda_q)$* , *Czechosl. Math. J.* **54** (129) (2004), 253–259.
18. R. Sahin, S. Ikikardes and Ö. Koruoğlu, *Generalized M^* -groups*, *Int. J. Algebra Comput.* **16** (2006), 1211–1219.
19. ———, *Some normal subgroups of the extended Hecke groups $\overline{H}(\lambda_p)$* , *Rocky Mountain J. Math.* **36** (2006), 1033–1048.
20. ———, *Extended Hecke groups $\overline{H}(\lambda_q)$ and their fundamental regions*, *Adv. Stud. Contemp. Math.* **15** (2007), 87–94.

BALIKESIR UNIVERSITY, DEPARTMENT OF MATHEMATICS, 10145, BALIKESIR, TURKEY

Email address: skardes@balikesir.edu.tr

BALIKESIR UNIVERSITY, DEPARTMENT OF MATHEMATICS, 10145, BALIKESIR, TURKEY

Email address: rsahin@balikesir.edu.tr