

# Estimation of a selected population parameters for Turkish marine waters' red mullet *Mullus barbatus ponticus* Essipov, 1927 (Actinopterygii: Perciformes: Mullidae)

## Estimación de parámetros poblacionales seleccionados para salmonetes de aguas marinas turcas *Mullus barbatus ponticus* Essipov, 1927 (Actinopterygii: Perciformes: Mullidae)

Dilek Türker 

Balikesir University, Department of Biology, Balikesir, Türkiye.  
Correspondence Author: [dturker@balikesir.edu.tr](mailto:dturker@balikesir.edu.tr)

### ABSTRACT

In order to calculate the length-weight and length-to-length relationships for red mullet *Mullus barbatus ponticus* Essipov, 1927 in Turkish marine waters, samples were collected from the Marmara Sea, the Northern Aegean Sea and the Western Black Sea by trawler from 2013 through 2016 fishing season. In the study in which 1,756 individuals were sampled, the number and proportion of individuals obtained from the Western Black Sea Region was 9.7% with 171 individuals from Amasra and 65.3% with 1,148 individuals from Ereğli; it was 1.7% with 30 individuals from the Marmara Sea, while it was 23.2% with 407 individuals from the North Aegean Sea. The total length was 6.3 to 20.5 cm, while the weight ranged from 2.54 to 89.53 g. In addition, 49 length-weight relationship values were examined from the literature of red mullet sampled in Turkish marine waters from 1977 to 2017. The smallest  $b$  value was determined as 2.84 from Iskenderun Bay and the highest was 3.3916 from the Izmir Bay. The median value of  $b$  was 3.188; it was observed that 50% of the  $b$  values ranged from 3.092 to 3.26.

**Key words:** *Mullus barbatus ponticus*; length-weight relationships; length-length relationships; growth parameters; positive allometry

### RESUMEN

Para calcular las relaciones longitud-peso y longitud-longitud de los salmonetes *Mullus barbatus ponticus* Essipov, 1927 distribuido en aguas del mar turco, se tomaron muestras del mar de Mármara, del mar Egeo septentrional y del mar Negro occidental mediante una red barredera durante el periodo de pesca de los años 2013 a 2016. En el estudio se tomaron muestras de 1.756 individuos. El número y la tasa de ejemplos obtenidos de la región occidental del Mar Negro, según las estaciones, fue del 9,7% con 171 individuos de Amasra y del 65,3% con 1.148 individuos de Ereğli; mientras que el 1,7% con 30 individuos del mar de Mármara, es del 23,2% con 407 individuos del mar Egeo septentrional. La altura total osciló de 6,3 a 20,5 cm, y el peso de 2,54 a 89,53 g. Además, se examinaron 49 artículos que publicaron datos de la relación longitud-peso de los salmonetes mostrados en aguas del mar turco de 1977 a 2017. Se encontró que el valor  $b$  más bajo fue de 2,84 en la Bahía de Iskenderun y el más alto de 3,3916 en la Bahía de İzmir. El valor promedio de  $b$  es 3,188; Se observó que el 50% de los valores  $b$  estaban en el intervalo de 3,092 a 3,26.

**Palabras clave:** *Mullus barbatus ponticus*; relación longitud-peso; relación longitud-longitud; parámetros de crecimiento; alometría positiva

## INTRODUCTION

According to fishbase [1], the goatfish family (Mullidae) includes 88 species belonging to 6 genera Worldwide. However, in Turkish marina waters the family Mullidae [2] is represented by *Mullus barbatus* Linnaeus, 1758, now *Mullus barbatus ponticus*, Essipov, 1927; *Mullus surmuletus* Linnaeus, 1758; *Parupeneus forsskali* (Fourmanoir & Guézé, 1976); *Upeneus moluccensis* (Bleeker, 1855) and *Upeneus pori* (Ben-Tuvia & Golani, 1989).

The red mullet (*M. barbatus* Linnaeus, 1758) is found in the Eastern Atlantic, including the Mediterranean and the Black Sea, from Western Norway, the English Channel (rare in the North Sea), to Dakar, Senegal, and the Canary Islands [3]. Red mullet's scientific name was *Mullus barbatus* Essipov, 1927, but fishbase changed it to *Mullus barbatus ponticus* Linnaeus, 1758 [1]. Red mullet is a significant target species for Mediterranean fisheries because of its commercial value [4, 5]. As a result, numerous writers have looked at the red mullet's population dynamics and biological characteristics in the Mediterranean [6, 7, 8, 9, 10, 11, 12, 13]. Fish biology, physiology, ecology, and fisheries evaluation all heavily rely on the length-weight relationships (LWRs) and length-length correlations (LLRs) [14]. They are applied in the assessment of fish stocks and populations [14, 15], and they are useful for between-region comparisons of life histories of species and the general health of fishing species, conditions, and reproduction history [16, 17, 18, 19, 20, 21, 22, 23, 24]. The Sea of Marmara, Western Black Sea, and Northern Aegean Sea are considered Turkey's most significant fishing grounds, contributing a sizeable share of the country's total marine fish production [25]. The goal of the current study was to examine the length-weight and length-length connections of red mullet samples collected from Turkey's Black Sea, Northern Aegean Sea, and Sea of Marmara coasts. (FIG. 1). Additionally, a total of 31 research conducted from 1977 to 2017 in Turkish marine waters yielded 49 LWRs. The following study evaluates Turkish scientific studies from the past and present.

## MATERIAL AND METHODS

During the 2012–2013 fishing seasons, samples of red mullet were taken using a commercial trawl net between the depths of 20–90 meters in the Sea of Marmara, Western Black Sea (Amasra and Ereğli sites), and Northern Aegean Sea. (FIG 1). The weights were recorded with a digital balance (AND GF 6100 Model, Japan) to the nearest 0.01 g, while the total length (TL), fork length (FL), and standard length (SL) were measured with digital calipers (Mitutoyo 500–181–30 Digital Compass, Japan) to the nearest mm. The equation  $W = aL^b$  [26, 27] was used to determine the length-weight relationships (LWRs), where  $W$  is the fish's body weight (g) and  $L$  is the entire length of the fish (cm). Least-squares regression was used to determine the parameters  $a$  and  $b$  as well as the coefficient of determination ( $r^2$ ). Additionally, 95% confidence limits of the parameter  $b$  were estimated by the equation:  $t = \left( \frac{SdlogTL}{SdlogW} \right) \times \frac{1b-3}{\sqrt{1-r^2}} \times (\sqrt{n-2})$ , where  $SdlogTL$  is the standard deviation of the logTL values,  $SdlogW$  is the standard deviation of the logW values,  $n$  is the number of samples used in the computation. If the estimated  $t$  value is greater than the tabular  $t$  values for  $n-2$  degrees of freedom, then the value of  $b$  differs from  $b = 3$  [27]. To determine whether parameter  $b$  and its confidence interval ( $= 0.05$ ) covers 3 or is significantly different from 3, the student's  $t$ -test was used to determine the growth type. Froese [28] found that changing length measurements had an effect on  $a$  but not  $b$ . Of particular note is the fact that for the same sample,  $a$  rise from total to fork and total to standard length [28]. Additionally, using linear regression analysis for TL–FL, FL–SL, and SL–TL, respectively, length-length relationships (LLRs) for the samples from the Sea of Marmara and Northern Aegean Sea were established.

## RESULTS AND DISCUSSION

A total of 1756 individuals were collected from Black Sea (9.7%;  $n=171$  from Amasra and 65.3%,  $n=1148$  from Ereğli), Sea of Marmara (1.7%,  $n=30$ ) and Northern Aegean Sea (23.2%,  $n=407$ ). The range of recorded total fish lengths and weights, parameters of LWRs and  $t$ -test results were calculated separately for each studied area and are presented in TABLE I.

Red mullet showed negative allometric growth ( $b < 3$ ) for Western Black Sea (Amasra) and Northern Aegean Sea (Edremit Bay). However, red mullet exhibited positive allometric growth ( $b > 3$ ) for Western Black Sea (Ereğli) and Sea of Marmara. The parameters of LLRs were given in TABLE II.

Length and weight characteristics, sex, number of individuals, values of  $a$ ,  $b$ ,  $r^2$  and sampling location of previous studies are given in TABLE III. The value of the parameter  $b$  in LWRs ranged from 2.84 in İskenderun Bay to 3.361 in Gökçeada Island for total length. The value of the parameter  $b$  in LWRs ranged from 2.9231 in İzmir Bay to 3.3916 in İzmir Bay for fork length [29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55].

According to Stergio and Moutopoulos [56] and Froese [28] a plot of  $\log(a)$  versus  $b$  for all known LWRs of a species results in a linear relationship which can be used to identify outliers. This method was applied for red mullet and the plots of  $\log(a)$  versus  $b$  for all available length-weight relationships (for each length type separately) are shown in FIG 2, 3, and 4.

There are various studies providing information about maximum length and also maximum weight of the species in coastal waters of Turkey (TABLE II). Results of the present study showed that minimum




FIGURE 1. Sampling areas (A: Edremit Bay (Northern Aegean Sea) B: Bandırma, Sea of Marmara; C: Ereğli and Amasra (Western Black Sea))

**TABLE I**  
Parameters of length-weight for red mullet

Location	Sex	n	Total Length (cm)		Weight (g)		Relationship Parameters			t-test	t-table (0.05)	GT
			Min	Max	Min	Max	r <sup>2</sup>	a	b	p		
Black Sea (Amasra)	C	171	6.3	15.0	2.54	31.75	0.918	0.0097	2.9941	0.0056	1.98	-A
Black Sea (Ereğli)	C	1148	6.8	16.1	2.60	40.79	0.9275	0.0063	3.19	0.1074	1.96	+A
Aegean Sea (Edremit)	C	407	10.1	20.5	8.45	89.53	0.8157	0.0157	2.795	0.0548	1.97	-A
Sea of Marmara (Bandırma)	C	30	11.1	18.5	12.50	62.86	0.9791	0.0067	3.1408	0.0463	2.04	+A

n: Sample size, Min: Minimum, Max: Maximum, a and b, Intercept and Slope of Length-Weight Relationships, r<sup>2</sup>: Coefficient of Determination, C, Combined, GT: Growth Type, -A: Negative Allometric, +A: Positive Allometric

**TABLE II**  
Parameters of Length-Length Relationships for red mullet

Location	TL (cm)			FL (cm)			SL (cm)			Equation	n	a	b	r <sup>2</sup>
	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max					
Northern Aegean Sea										TL = a + bFL		0.2460	1.0821	0.9486
Northern Aegean Sea	12.66	10.1	20.5	11.47	9.2	18.8	10.08	8.0	16.6	FL = a + bSL	407	0.7321	1.0649	0.9482
Northern Aegean Sea										SL = a + bTL		-0.0724	0.8023	0.9505
Sea of Marmara										TL = a + bFL		0.5311	1.0824	0.9855
Sea of Marmara	14.26	11.1	18.5	12.68	9.5	16.5	11.24	8.5	14.4	FL = a + bSL	30	-0.2618	1.1514	0.9834
Sea of Marmara										SL = a + bTL		0.0236	0.7868	0.9921

TL: Total Length, FL: Fork Length, SL: Standard Length, Min: Minimum, Max: Maximum, n: Sample size, a and b: intercept and slope of Length-Length Relationship, r<sup>2</sup>: coefficient of determination

**TABLE III**  
49 Length-Weight Relationships Obtained from Different Areas

Location	L min-L max (cm)	LT	W min-W max (g)	Sex	n	a	b	r <sup>2</sup>	Reference
İzmir Bay	7.50-22.0	FL	6.00-115.00	C	6054	0.0165	2.9231	-	[29] Toğulga, 1977
İzmir Bay	8.60-18.3	FL	9.00-121.00	F	218	0.0059	3.3916	0.9890	[30] Akyol et al., 2000
İzmir Bay	9.50-15.0	FL	12.00-51.00	M	110	0.0077	3.2834	0.9750	[30] Akyol et al., 2000
İzmir Bay	8.60-18.3	FL	9.00-121.00	C	346	0.0063	3.3625	0.9880	[30] Akyol et al., 2000
Edremit Bay	9.45-18.7	FL	13.45-87.65	C	474	0.0157	2.9811	0.9620	[31] Çelik and Torcu, 2000
İzmir Bay	-	FL	-	F	155	0.0073	3.2800	-	[11] Kınacıgil et al., 2001
İzmir Bay	-	FL	-	M	65	0.0077	3.2500	-	[11] Kınacıgil et al., 2001
İzmir Bay	8.10-16.1	FL	7.00-66.00	C	220	0.0071	3.2900	-	[11] Kınacıgil et al., 2001
Babadillimani Bight	3.80-21.5	TL	0.39-119.90	C	2021	0.0076	3.1280	0.9760	[32] Çiçek et al., 2006
SE Black Sea	6.80-18.0	TL	-	F	248	0.0047	3.2700	0.9800	[33] Demirhan & Can, 2006
SE Black Sea	6.80-14.6	TL	-	M	173	0.0057	3.1900	0.9400	[33] Demirhan & Can, 2006
SE Black Sea	6.80-14.6	TL	-	C	432	0.0051	3.2400	0.9700	[33] Demirhan & Can, 2006
N. Aegean Sea	12.70-22.3	TL	-	F	49	0.0038	3.3610	0.9350	[34] Karakulak et al., 2006
N. Aegean Sea	12.50-18.6	TL	-	M	16	0.0067	3.1710	0.9420	[34] Karakulak et al., 2006
N. Aegean Sea	12.50-22.3	TL	-	C	76	0.0049	3.2730	0.9410	[34] Karakulak et al., 2006
İzmir Bay	7.50-20.0	FL	5.57-123.00	C	479	0.0102	3.1760	0.9600	[35] Özaydın&Taşkavak,06
İzmir Bay	8.00-19.6	TL	6.00-90.00	C	111	0.0091	3.1000	0.9700	[36] Gökçe et al., 2007
Saros Bay	6.00-24.7	TL	2.00-200.00	C	3386	0.00762	3.0949	0.9630	[37] İşmen et al., 2007
Middle Black Sea	6.60-18.4	TL	2.94-60.16		176	0.0111	2.9630	0.9800	[38] Kalaycı et al., 2007
İzmir Bay	5.40-21.2	FL	-	C	1910	0.0089	3.2330	0.9810	[39] Özaydın et al., 2007

**TABLE III cont...**  
**49 Length-Weight Relationships Obtained from Different Areas**

NE Mediterranean	8.20-22.0	TL	4.96-106.26	C	451	0.0032	3.0600	0.9400	[40] Sangun <i>et al.</i> , 2007
İzmir Bay	8.20-28.2	TL	-	F	970	0.0056	3.2400	0.9800	[41] İlkyaz <i>et al.</i> , 2008
İzmir Bay	8.20-19.0	TL	-	M	909	0.0064	3.1900	0.9640	[41] İlkyaz <i>et al.</i> , 2008
İzmir Bay	8.20-28.20	TL	-	C	1879	0.0060	3.2200	0.9780	[41] İlkyaz <i>et al.</i> , 2008
İskenderun Bay	11.00-20.40	TL	15.98-91.30	C	8	0.0184	2.8400	0.9900	[42] Gökçe <i>et al.</i> , 2010
Çandarlı Bay	4.60-9.90	TL	0.55-8.01	C	13	0.0040	3.3440	0.9540	[43] Gürkan <i>et al.</i> , 2010
N. Sea of Marmara	10.00-15.70	TL	9.54-46.59	C	99	0.0049	3.3260	0.9160	[44] Bök <i>et al.</i> , 2011
Sea of Marmara	9.60-22.70	TL	-	C	94	0.0150	3.0040	0.8600	[45] Demirel&Dalkara,2012
Gallipoli Peninsula and Dardanelles	8.70-20.10	TL	6.83-99.13	C	102	0.0062	3.2200	0.9800	[46] Cengiz, 2013
HomaLagoon,İzmir	5.10-11.10	TL	1.15-13.82	C	90	0.0060	3.1800	0.9910	[47] Acarlı <i>et al.</i> , 2014
Saros Bay	9.20-23.60	TL	7.50-177.30	F	2302	0.0610	3.1900	0.9400	[48] Arslan & İşmen, 2014
Saros Bay	8.80-24.10	TL	7.80-119.70	M	1308	0.0800	3.0890	0.9200	[48] Arslan & İşmen, 2014
Saros Bay	6.50-24.80	TL	2.50-177.30	C	9386	0.0084	3.0770	0.9400	[48] Arslan & İşmen, 2014
S. Aegean Sea	5.60-38.20	TL	-	C	2009	0.0065	3.3550	0.9700	[49] Bilge <i>et al.</i> , 2014
Gulf of Antalya	8.70-21.50	TL	-	C	1565	0.0071	3.1650	0.8940	[50] Özvarol, 2014
Black Sea	5.30-19.00	TL	1.20-73.40		2693	0.0074	3.1230	0.9600	[51] Kasapoğlu & Düzgüneş 2014
Çandarlı Bay	5.20-22.40	FL	1.50-146.10	C	970	0.0064	3.3340	0.9890	[52] Akalın <i>et al.</i> , 2015
İskenderun Bay	6.90-15.70	TL	3.41-51.38	C	212	0.0072	3.1618	0.9530	[12] Çiçek, 2015
Eastern Black Sea	11.10-21.40	TL	12.41-96.22	F	433	0.0064	3.1340	0.9240	[53] Yeşilçiçek <i>et al.</i> , 2015
Eastern Black Sea	9.40-19.80	TL	8.49-66.21	M	212	0.0090	2.9930	0.8890	[53] Yeşilçiçek <i>et al.</i> , 2015
Eastern Black Sea	7.40-22.60	TL	2.68-102.50	C	672	0.0066	3.1190	0.9250	[53] Yeşilçiçek <i>et al.</i> , 2015
İzmir Bay	9.50-13.40	TL	9.00-27.10	C	47	0.0068	3.1930	0.9997	[54] Kara <i>et al.</i> , 2016
Western Black Sea	9.10-18.90	TL	8.81-62.42	F	1986	0.0103	3.0127	0.9407	[13] Yıldız & Karakulak, 2016
Western Black Sea	9.00-15.80	TL	7.18-47.97	M	1829	0.0137	2.8993	0.9283	[13] Yıldız & Karakulak, 2016
Western Black Sea	6.30-18.90	TL	3.62-62.42	C	4928	0.0109	2.9886	0.9554	[13] Yıldız & Karakulak, 2016
İzmir Bay	4.50-11.90	TL	0.75-16.80	C	107	0.0062	3.1900	0.9970	[54] Kara <i>et al.</i> , 2016
Gülbağçe Bay	-	FL	-	F	301	0.0113	3.1520	0.9660	[55] Kurtul & Özyayın, 2017
Gülbağçe Bay	-	FL	-	M	229	0.0102	3.1880	0.9590	[55] Kurtul & Özyayın, 2017
Gülbağçe Bay	5.10-15.30	FL	-	C	629	0.0100	3.2010	0.9720	[55] Kurtul & Özyayın, 2017

LT: Length Type, C: Combined, F: Female, M: Male, TL: Total Length, FL: Fork Length, n: Sample Size,  $a$  and  $b$  Intercept and Slope of Length-Weight Relationships,  $r^2$ : Coefficient of Determination) (SE: Southeastern, N: Northern, S: Southern




FIGURE 2. The log ( $a$ ) vs  $b$  of red mullet for Fork Length





FIGURE 3. The log ( $a$ ) vs  $b$  of red mullet for Total Length. The circled points are outliers



**FIGURE 4.** Scatter plot of mean  $\log(a)$  over mean  $b$  for red mullet. Areas of negative allometric, isometric, positive allometric change in body weight relative to body length are indicated

total length of studied locations was longer than some previous studies, but the maximum total length of studied locations was shorter than some previous studies (TABLE I). There may be many reasons for this situation, but the ecological conditions of the fish sampled from stations representing three different marine environments (Black Sea, Marmara Sea and Aegean Sea) are perhaps the most important reason. In addition, it is thought that the use of different fishing gear, due to the characteristics of these three different marine environments and the decisions of policy makers, causes different results.

The red mullet LWRs'  $b$  values from this study are comparable to those from earlier research by Toğulga [29] and Kurtul and Zaydn [55]. The habitat, region, seasonal effects, degree of stomach fullness, gonad maturity, sex, health, preservation methods, and variations in the observed length ranges of the samples caught may all have an impact on fish LWRs [20, 39, 57]. However, all of these factors were taken into consideration in this study. If there are numerous studies of LWRs for a species, outliers can be found using a scatter plot of  $\log(a)$  and  $b$  values, according to Stergiou and Moutopoulos and Froese [28, 56]. As a result, 31 studies on red mullet from Turkish maritime waters from 1977 to 2017 were examined. (FIG 2 and 3). Problematic studies for red mullet are shown by the circled outliers. Froese [28] claims that a stronger regression analysis conducted after removing outlier observations would be powerful enough to explain the %99 of the variance [28]. These outliers also result in a decline. FIG 4 demonstrates that several of the estimations were higher than 3 when a scatter plot for  $\log(a)$  and  $b$  was produced. This enables the conclusion to be drawn that red mullet found in Turkish waters primarily exhibit positive allometric growth in both fork and total length.

## CONCLUSIONS AND IMPLICATIONS

This study presents LWRs and LLRs from Sea of Marmara, Western Black Sea and Northern Aegean Sea and a collected list of the LWRs parameters for red mullet in Turkish marine waters from previous published studies. These important data and results may be used by fishery management authorities and further academic studies. Therefore, relevant studies could be supportive in future for the management of red mullet fisheries in Turkey. Additionally, updating

information about the maximum size of a species that might be commercially or recreationally exploited in the future is important.

## Conflicts of interest

The authors declare that there is no conflict of interest in this work.

## BIBLIOGRAPHIC REFERENCES

- [1] Froese R, Pauly D, editors. FishBase [Internet]. Version 10/2023. Paris: FishBase Team. 2023 [cited 18 Oct. 2023]. Available in: <http://www.fishbase.org>.
- [2] Bilecenoğlu M, Taşkavak E, Mater S, Kaya M. Checklist of the marine fishes of Turkey. Zootaxa [Internet]. Auckland, New Zealand: Magnolia Press. 2002; 113(1):1-194. doi: <https://doi.org/mbr9>
- [3] Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J, Tortonese E, editors. Fishes of the north-eastern Atlantic and the Mediterranean. Vol. 3. Paris: Unesco; 1986; p. 511-1473.
- [4] Demestre M, Sbrana M, Álvarez F, Sánchez P. Analysis of the interaction of fishing gear in *Mullus barbatus* fisheries of the Western Mediterranean. J. Appl. Ichthyol. [Internet]. 1997; 13(2):49-56. doi: <https://doi.org/dt99mx>
- [5] Tserpes G, Fiorentino F, Levi D, Cau A, Murenu M, Zamboni A, Papaconstantinou C. Distribution of *Mullus barbatus* and *M. surmuletus* (Osteichthyes: Perciformes) in the Mediterranean continental shelf: implications for management. Scient. Marina. [Internet]. 2002; 66(S2):39-54. doi: <https://doi.org/dqfc7h>
- [6] Livadas RJ. A study of the growth and maturity of striped mullet (*Mullus barbatus* L.) in waters of Cyprus. Rome: Food and Agriculture Organization of the United Nations. 1988. FAO Fisheries Report No. 412. p. 44-51.
- [7] Vrantzas N, Kalagia M, Karlou C. Age, growth and state of stock of red mullet (*Mullus barbatus* L., 1758) in the Saronikos Gulf of Greece. Rome: Food and Agriculture Organization of the United Nations. 1992. FAO Fisheries Rep No. 477 p. 51-67.



- [8] Levi D, Andreoli G, Giusto GB. An analysis based on trawl-survey data of the state of the "Italian" stock of *Mullus barbatus* in the Sicilian Channel, including management advice. Fisheries Res. [Internet]. 1993; 17:334–341. doi: <https://doi.org/c75svh>
- [9] Tursi A, Matarrese A, D'Onghia G, Sion L. Population biology of red mullet (*Mullus barbatus* L.) from the Ionian Sea. Marine Life. 1994; 4(2):33–43.
- [10] Fiorentino F, Zamboni A, Rossi M, Relini G. The growth of the red mullet (*Mullus barbatus*, L. 1758) during the first years of life in the Ligurian Sea (Mediterranean). Cahiers Options Mediterranennes. [Internet]. 1998 [cited 24 Sept 2023]; (35):65–78. Available in: <https://bit.ly/3TSuKpo>.
- [11] Kinacigil HT, İlkyaz AT, Akyol O, Metin G., Çıra E, Ayaz A. Growth parameters of red mullet (*Mullus barbatus* L., 1758) and seasonal cod-end selectivity of traditional bottom trawl nets in İzmir Bay (Aegean Sea). Acta Adriatica. [Internet]. 2001 [cited 03 Oct 2023]; 42(1):113–123. Available in: <https://bit.ly/3SbxPj7>.
- [12] Çiçek E. Age, growth and mortality parameters of *Mullus barbatus* Linnaeus, 1758 (Perciformes: Mullidae) in İskenderun Bay, northeastern Mediterranean. Iranian J. Ichthyol. 2015; 2(4):262–269.
- [13] Yıldız T, Karakulak FS. An investigation of age, growth and mortality of the red mullet *Mullus barbatus* Linnaeus, 1758 in the western Black Sea. Cahiers de Biologie Marine. 2016; 57(4):415–425.
- [14] Gonçalves JMS, Bentes L, Lino PG, Ribeiro J, Canario AV, Erzini K. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west Coast of Portugal. Fisheries Res. [Internet]. 1997; 30(3):253–256. doi: <https://doi.org/d9m3jv>
- [15] Ricker WE. Handbook of computations for biological statistics of fish populations. Ottawa, Canada: Fisheries Research Board of Canada. 1958; Series Bulletin No. 119. 300 p.
- [16] Petrakis G, Stergiou KI. Weight-length relationships for 33 fish species in Greek Waters. Fisheries Res. [Internet]. 1995; 21(3–4):465–469. doi: <https://doi.org/cbm7qk>
- [17] Kara A, Bayhan B. Length-weight and length-length relationships of the bogue *Boops boops* (Linnaeus, 1758) in İzmir Bay (Aegean Sea of Turkey). Belgian J. Zool. 2008; 138(2):154–157.
- [18] Nikolsky GV. The ecology of fishes. London, UK: Academic Press; 1963. 352 p.
- [19] Wootton RJ. The evolution of life histories: theory and analysis. Rev. Fish Biol. Fisheries. [Internet]. 1993; 3(4):384–385. doi: <https://doi.org/cdvp5k>
- [20] Wootton RJ. Ecology of teleost fishes. Dordrecht, Netherlands: Kluwer Academic Publishers; 1998. 403 p.
- [21] Pauly D. Fishbyte Section. Editorial. Naga, The ICLARM Quarterly. 1993. 16:26.
- [22] Erkoyuncu İ. Balıkçılık biyolojisi ve populasyon dinamiği [Fisheries Biology And Population Dynamics]. Sinop, Türkiye: Ondokuz Mayıs Üniversitesi Yayınları; 1995; 265 p. Turkish.
- [23] Aşar D. Balıkçılık biyolojisi ve populasyon dinamiği [Fisheries Biology And Population Dynamics]. Adana, Türkiye: Nobel Kitabevi; 2005; 288 p. Turkish.
- [24] Radkhan A, Eagderi S. Length-weight and length-length relationships and condition factor of six cyprinid fish species of Zarrineh River (Urmia Lake Basin, Iran). Iranian J. Ichthyol. [Internet]. 2015 [cited 15 Sept 2023]; 2(1):61–64. Available in: <https://bit.ly/3RSPx9F>.
- [25] Fishery statistics, State Institute of Statistics. Prime Minister Republic of Turkey. 2018; 3 p.
- [26] Ricker WE. Computing and interpretation of biological statistics of fish populations. Ottawa, Canada: Fisheries Research Board of Canada. 1975; Series Bulletin No 191: 382 p.
- [27] Pauly D. Fish population dynamics in tropical waters: a manual for use with programmable calculators. Manila, Philippines: International Center for Living Aquatic Resources Management (ICLARM). ICLARM Studies Rev. 8; 1984. 325 p.
- [28] Froese R. Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. J. Appl. Ichthyol. [Internet]. 2006; 22(4):241–253. doi: <https://doi.org/db6sns>
- [29] Toğulga M. The studies on population dynamics of red mullet (*Mullus barbatus* Lin. 1758) in İzmir Bay. Ege Univ. J. Faculty Sci. 1977; 2:175–194.
- [30] Akyol O, Tosunoğlu Z, Tokaç A. Investigations of the growth and reproduction of red mullet (*Mullus barbatus* Linnaeus, 1758) population in the Bay of İzmir (Aegean Sea). Anadolu Univ. J. Sci. Technol. 2000; 1(1):121–127.
- [31] Çelik O, Torcu H. Investigations on the biology of red mullet (*Mullus barbatus* Linnaeus, 1758) in Edremit Bay, Aegean Sea, Turkey. Turkish J. Vet. Anim. Sci. 2000; 24(3):287–295.
- [32] Çiçek E, Aşar D, Yeldan H, Özütok M. Length-weight relationships for 31 teleost fishes caught by bottom trawl net in the Babadillmani Bight (northeastern Mediterranean). J. Appl. Ichthyol. [Internet]. 2006; 22(4):290–292. doi: <https://doi.org/d6t8z8>
- [33] Demirhan SA, Can MF. Length-weight relationships for seven fish species from the southeastern Black Sea. J. Appl. Ichthyol. [Internet]. 2007; 23(3):282–283. doi: <https://doi.org/ccgpr6>
- [34] Karakulak FS, Erk H, Bilgin B. Length-weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. J. Appl. Ichthyol. [Internet]. 2006; 22(4):274–278. doi: <https://doi.org/bpfn8j>
- [35] Özyayın O, Taşkavak E. Length-weight relationships for 47 fish species from İzmir Bay (eastern Aegean Sea, Turkey). Acta Adriatica. 2006; 47(2):211–216.
- [36] Gökçe G, Aydın I, Metin C. Length-weight relationships of 7 fish species from the north Aegean Sea, Turkey. Intern. J. Natural Engineering Sci. 2007; 1:51–52.
- [37] İşmen A, Özen O, Altınağaç U, Özekinci U, Ayaz A. Weight-length relationships of 63 fish species in Saros Bay, Turkey. J. Appl. Ichthyol. [Internet]. 2007; 23(6):707–708. doi: <https://doi.org/dqj3b>
- [38] Kalaycı F, Samsun N, Bilgin S, Samsun O. Length-weight relationship of 10 fish species caught by bottom trawl and midwater trawl from the Middle Black Sea, Turkey. Turkish J. Fisheries Aquatic Sci. [Internet]. 2007 [cited 14 June 2023]; 7(1):33–36. Available in: <https://bit.ly/3Hbn0qU>.

- [39] Özyayın O, Uçkun D, Akalın S, Leblebici S, Tosunoğlu Z. Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *J. Appl. Ichthyol.* [Internet]. 2007; 23(6):695-696. doi: <https://doi.org/cjbb2b>
- [40] Sangun L, Akamca E, Akar M. Weight-length relationships for 39 fish species from the north-eastern Mediterranean Coast of Turkey. *Turkish J. Fisheries Aquatic Sci.* [Internet]. 2007 [cited 14 June 2023]; 7(1):37-40. Available in: <https://bit.ly/3HdBczH>.
- [41] İlkayaz AT, Metin G, Soykan O, Kinacıgil HT. Length-weight relationships of 62 fish species from the Central Aegean Sea, Turkey *J. Appl. Ichthyol.* [Internet]. 2008; 24(6):699-702. doi: <https://doi.org/fbp385>
- [42] Gökçe G, Çekiç M, Filiz H. Length-weight relationships of marine fishes of Yumurtalık Coast (İskenderun Bay), Turkey. *Turkish J. Zool.* [Internet]. 2010; 34(1):101-104. doi: <https://doi.org/mbv5>
- [43] Gürkan S, Bayhan B, Akçınar SC, Taşkavak E. Length-weight relationship of fish from Shallow Waters of Çandarlı Bay (north Aegean Sea, Turkey). *Pakistan J. Zool.* [Internet]. 2010 [cited 18 June 2023]; 42(4):495-498. Available in: <https://bit.ly/48MIANZ>.
- [44] Bök TD, Göktürk D, Kahraman AE, Aliçlı TZ, Acun T, Ateş C. Length-weight relationships of 34 fish species from the Sea of Marmara, Turkey. *J. Anim. Vet. Adv.* [Internet]. 2011 [cited 18 June 2023]; 10(23):3037-3042. Available in: <https://bit.ly/48oAcER>.
- [45] Demirel N, Dalkara-Murat E. Weight-length relationships of 28 fish species in the Sea of Marmara. *Turkish J. Zool.* [Internet]. 2012; 36(6):785-791. doi: <https://doi.org/mbv6>
- [46] Cengiz Ö. Length-weight relationships of 22 fish species from The Gallipoli Peninsula and Dardanelles (Northeastern Mediterranean, Turkey). *Turkish J. Zool.* [Internet]. 2013; 37(4):419-422. <https://doi.org/mbv7>
- [47] Acarlı D, Kara A, Bayhan B. Length-weight relationships for 29 fish species from Homa Lagoon, Aegean Sea, Turkey. *Acta Ichthyol. et Piscatoria.* [Internet]. 2014; 44(3):249-257. doi: <https://doi.org/mbv8>
- [48] Arslan M, İşmen A. Age, growth, reproduction and feeding of *Mullus barbatus* in Saros Bay (North Aegean Sea). *Journal Black Sea/ Mediterranean Environ.* [Internet]. 2014 [cited 4 Aug. 2023]; 20(3):184-199. Available in: <https://bit.ly/3TXhkbL>.
- [49] Bilge G, Yapıcı S, Filiz H, Cerim H. Weight-length relations for 103 fish species from the southern Aegean Sea, Turkey. *Acta Ichthyol. et Piscatoria.* [Internet]. 2014; 44(3):263- 269. doi: <https://doi.org/mbv9>
- [50] Özvarol Y. Length-weight relationships of 14 fish species from the Gulf of Antalya (northeastern Mediterranean Sea, Turkey). *Turkish J. Zool.* [Internet]. 2014; 38(3):342-346. doi: <https://doi.org/mbwb>
- [51] Kasapoğlu N, Düzgüneş E. Length-weight relationships of marine species caught by five gears from the Black Sea. *Mediterranean Marine Sci.* [Internet]. 2014; 15(1):95-100. doi: <https://doi.org/mbwc>
- [52] Akalın S, İlhan D, Özyayın O. Length-weight relationships for 30 demersal fish species from Çandarlı Bay (north Aegean Sea, Turkey). *Croatian J. Fisheries.* [Internet]. 2015; 73(2):73-76. doi: <https://doi.org/mbwd>
- [53] Yeşilçiçek T, Kalaycı F, Şahin C. Length-weight relationships of 10 fish species from the southern Black Sea. *Turkey J. Fisheries Sci.* [Internet] 2015 [cited 25 Aug 2023]; 9(1):19-23. Available in: <https://bit.ly/4aM1E0Q>.
- [54] Kara A, Sağlam C, Acarlı D. Length-weight relationships of fish captured by wire pots in Izmir Bay (eastern Aegean Sea, Turkey). [Internet]. In: Chatziefstathiou M, editor. *Proceedings of the 2nd International Congress on Applied Ichthyology & Aquatic Environment*; 10 - 12 Nov 2016; Messolonghi, Greece. Volos, Greece: University of Thessaly. [cited 25 Oct 2023]. p. 250-253. Available in: <https://bit.ly/3TS0Ame>.
- [55] Kurtul I, Özyayın O. Age, growth and length-weight relationship of red mullet (*Mullus barbatus* Linnaeus, 1758) in Gülbahçe Bay (Aegean Sea). *Turkish J. Aquatic Sci.* [Internet] 2017; 32(3):135-143. doi: <https://doi.org/mbwj>
- [56] Stergiou KI, Moutopoulos DK. A review of length-weight relationships of fishes from Greek Marine Waters. *Naga, The ICLARM Quarterly.* 2001; 24(1-2):23-39.
- [57] Tesch FW. Age and growth. In: Ricker WE. editor. *Methods for assessment of fish production in fresh waters.* Oxford, UK: Blackwell Scientific. 1971. p. 99-130.