

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/287613499>

Comparative analysis of isokinetic leg strength in professional soccer and basketball players

Article in *South African Journal for Research in Sport, Physical Education and Recreation* · January 2013

CITATIONS

3

READS

128

1 author:



Ibrahim Erdemir

Balıkesir University

16 PUBLICATIONS 32 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Effects of Water Intake on Biochemical Parameters and Performance During Resistance Exercise [View project](#)

COMPARATIVE ANALYSIS OF ISOKINETIC LEG STRENGTH IN PROFESSIONAL SOCCER AND BASKETBALL PLAYERS

Ibrahim ERDEMIR

School of Physical Education and Sport, Balikesir University, Balikesir, Turkey

ABSTRACT

This study aimed to compare and analyse flexion and extension peak torques and hamstring to quadriceps peak torque muscle ratio of the knee joints by measuring isokinetic knee strength in professional soccer and basketball players. Twenty-two players were recruited, which included 12 soccer players (age: 15.8±1.0 years) and 10 basketball players (age: 15.7±0.9 years). After recording the stature, body mass, body mass index, body fat percentage, body fat mass and vertical jump measurement of each participant, lower extremity knee-joint measurements were conducted using an isokinetic dynamometer at angular velocities of 60°·s⁻¹ and 240°·s⁻¹. Peak torques of the hamstring and quadriceps muscles at 60°·s⁻¹ and 240°·s⁻¹ were significantly higher in basketball than soccer players (p<0.05 to p<0.01). When the hamstring to quadriceps peak torque muscle ratios were compared between soccer and basketball players, the only significant difference was found in the left knee of basketball players at 60°·s⁻¹. However, the relative strength of the hamstring and quadriceps muscles did not differ between soccer and basketball players at 60°·s⁻¹ and 240°·s⁻¹. In conclusion, body mass has a decisive effect on the production of peak torque values of quadriceps and hamstring muscles in soccer and basketball players.

Key words: Hamstring; Quadriceps ratio; Peak torque; Knee flexion; Knee extension; Isokinetic strength.

INTRODUCTION

Muscle strength is one of the key factors in successful sport performance and is an important indicator of the effectiveness of injury rehabilitation in athletes (Croisier *et al.*, 2002). To monitor the performance of athletes, as well as the rehabilitation progress of injured players, various lower limb strength indices have been investigated. Among these, hamstring-to-quadriceps muscle peak torque muscle ratio (H:Q muscle ratio) is one of the most commonly evaluated. This ratio of strength of agonist to antagonist leg muscles has been used to examine functional ability, knee joint stability and muscle balance between hamstring (H) and quadriceps (Q) muscles during velocity-dependent movements (Aagaard *et al.*, 1995; Zakas *et al.*, 1995; Li *et al.*, 1996; Orchard *et al.*, 1997; Clanton & Coupe, 1998; Hewett *et al.*, 1999). An injury may occur during rapid leg extension if the hamstrings fail to generate effective eccentric counteraction to decelerate the movement (Croisier *et al.*, 2008). Further, when the hamstrings act to extend the hip, muscle strains may occur during rapid alterations between flexion and extension (Petersen & Holmich, 2005). The anterior cruciate ligament, assisted by the hamstring muscles, stabilises the knee by preventing anterior translation of the tibia on the femur (Kannus, 1988; Moore & Wade, 1989; Pettitt & Bryson, 2002), which can

occur during pivoting movements, such as landing from a jump and sudden changes in direction in field (soccer) and court (volleyball and basketball) athletes (Griffin *et al.*, 2000).

One of the most important motor skills in sport is referred to as the vertical jump (Barnett *et al.*, 2008). Thissen-Milder and Mayhew (1990) hypothesise that the jump is an important factor for most athletes in sport, because jumping is a major part of attack and defence movements in all sport games. The vertical jump is described as a ballistic movement and consists of rapid eccentric muscle activity followed by maximal concentric actions. Performing the motor movements requires the ability to strengthen the muscles and dynamic stabilisers of the knee. The extensors, dynamic knee stabiliser muscles based on the requirements of the sport, function mainly in the impulsive phase of the jump and the landing phase (Panni *et al.*, 2002).

The demands of certain sport on the knee joint may be related to the high incidence of injury in that joint (Richards *et al.*, 1996), and may cause imbalances in muscle strength between antagonistic dynamic knee stabiliser muscles. These muscle imbalances in muscle force production may increase the risk of injury because of the resultant high levels of stress in tissues (Oberge *et al.*, 1986). The isokinetic dynamometer provides fast and reliable quantification of variables related to muscle performance at different angular velocities, including the maximum torque, total work, reciprocity between agonist and antagonist muscles and fatigue index (Perrin *et al.*, 1987). Isokinetic assessment allows the identification of muscle strength deficit between bilateral muscle groups and between reciprocal muscle groups (agonist and antagonist) (Siqueira *et al.*, 2002).

There are several studies that used isokinetic dynamometer measurements in different populations (Kazazovic *et al.*, 2008). However, there is little information about the measurements used for athletes in team sport, especially in research of the lower extremities (Kazazovic *et al.*, 2009). Basketball and soccer are among the most popular sports. Both sports use the lower extremities and consist of some complex movements which require strenuous efforts, such as sudden feints, stops, starts, duels, sprints and jumps (Reilly & Thomas, 1976). These efforts depend on the strength of the neuromuscular system, especially in the lower limbs (Cometti *et al.*, 2001).

The aim of this study, therefore, was to compare and analyse the maximal voluntary peak torques of the quadriceps and hamstring muscles, and the torque ratio between these muscle groups of the right and left legs in professional basketball and soccer players by using isokinetic tests.

MATERIALS AND METHODS

Participants

The study included 12 male professional (division III) soccer players (age: 15.83±1.03 years; body mass: 59.25±6.43kg; stature: 171.50±4.89cm) and 10 male professional basketball players (division III) (age: 15.67±0.87 years; body mass: 75.68±13.27kg; stature: 185.67±8.93cm). All of the participants were fully informed of the goals and methodology of the test and provided signed consent. The participants agreed with the testing process and the

use of the data for further research. The day before testing, the players were not subjected to any intense training.

Prior to participation in the study, the players were interviewed about their medical records and completed an injury questionnaire. Participants were excluded from participation in the study if they had any current hip, knee, or ankle injury or any other leg injury. The participants were all right-leg dominant.

Physical measurements

Stature and body mass were measured with an electronic scale (708 Seca, Hamburg, Germany). Body mass index (BMI), body fat percentage and body fat mass were measured with a Tanita Body Composition Analyser BC-418, using the bioelectrical impedance analysis method. The vertical jump test was also performed using a specialised apparatus called the Vertec. Before the testing session started, the participants were allowed a 15-minute warm-up at a light intensity on the leg curl and leg extension machine.

Isokinetic measurements

Participants were tested in the sitting position on IsoMed 2000 isokinetic dynamometer. Participants were seated for testing in the chair of the dynamometer with the backrest angle at 90°. The axis of rotation of the right knee was aligned with the axis of rotation of the dynamometer's armature, and the ankle cuff was attached approximately 3cm above the dorsal surface of the foot. Gravity correction was performed before the test. Stabilisation straps were placed over the pelvis and chest, and participants positioned their arms across their chests during familiarisation and testing. To synchronise themselves with the testing device, participants were instructed to perform 3 active repetitions of knee movement ranging from maximal flexion to maximal extension.

To adapt to the test conditions, participants were allowed 3 sub-maximal contractions of the quadriceps and hamstring muscle groups at the beginning of the tests. Standardised verbal motivation techniques were used to encourage maximal work from the test participants. All participants performed 10 maximal (the first and the last of the 10 were dismissed) concentric contractions (knee flexion and extension) of both legs at velocities of 60°·s⁻¹ and 240°·s⁻¹ (Brockett *et al.*, 1999). A rest period of 3 minutes was allowed between test speeds, and 5 minutes were allowed between test limbs.

Statistical analysis

The data were processed with SPSS 15.0 for Windows (SPSS Inc., USA). Statistical parameters were calculated for all of the variables, and one-way analysis of variance (ANOVA) was applied to determine the statistically significant differences between basketball and soccer players. Statistical significance was set at the level of $p < 0.05$.

RESULTS

Age and physical characteristics of the participants are presented in Table 1. No significant between-group differences in age, BMI and vertical jump were noted. However, the mean of

stature, body mass and body fat mass were significantly ($p<0.01$) lower in soccer players than basketball players. Further, body fat percentage was also significantly ($p<0.05$) lower in soccer players than basketball players (Table 1).

TABLE 1: MEAN \pm SD FOR AGE AND PHYSICAL CHARACTERISTICS OF PARTICIPANTS

Parameter	Soccer (n = 12)	Basketball (n = 10)
Age (years)	15.8 \pm 1.03	15.7 \pm 0.9
Stature (cm)	171.5 \pm 4.9	185.7 \pm 8.9**
Body mass (kg)	59.3 \pm 6.4	75.7 \pm 13.3**
Body mass index (kg/m ²)	20.1 \pm 1.8	22.1 \pm 4.3
Body fat (%)	11.0 \pm 4.5	15.5 \pm 5.1*
Body fat mass (kg)	6.6 \pm 2.8	12.1 \pm 5.9**
Vertical jump (cm)	42.8 \pm 3.2	48.4 \pm 7.9

Significant differences: * $p < 0.05$; ** $p < 0.01$

TABLE 2: MEAN VALUES (\pm SD) OF HAMSTRING AND QUADRICEPS PEAK TORQUES AT VELOCITIES OF 60°S⁻¹ AND 240°S⁻¹

Variable	Angular Velocity LEFT		Angular Velocity RIGHT	
	Soccer (n=12)	Basketball (n=10)	Soccer (n=12)	Basketball (n=10)
PT Hamstring (Nm)				
60°s ⁻¹	96.8 \pm 16.5	138.0 \pm 24.8**	103.8 \pm 10.0	138.8 \pm 27.1**
240°s ⁻¹	82.4 \pm 21.2	115.1 \pm 27.8*	85.7 \pm 12.0	118.7 \pm 28.9**
PT Quadriceps (Nm)				
60°s ⁻¹	203.3 \pm 33.4	244.0 \pm 37.2*	199.2 \pm 33.0	252.8 \pm 40.5**
240°s ⁻¹	125.3 \pm 17.0	168.6 \pm 40.3*	123.6 \pm 24.7	168.7 \pm 31.2**
PT Hamstring: Quadriceps (%)				
60°s ⁻¹	48.4 \pm 7.7	56.7 \pm 4.8*	53.2 \pm 7.0	55.1 \pm 6.0
240°s ⁻¹	66.3 \pm 16.5	69.0 \pm 9.9	70.8 \pm 10.3	70.9 \pm 7.2
PT Hamstring/Body Mass				
60°s ⁻¹	1.6 \pm 0.2	1.8 \pm 0.3	1.8 \pm 0.2	1.9 \pm 0.3
240°s ⁻¹	1.4 \pm 0.4	1.5 \pm 0.4	1.45 \pm 0.2	1.6 \pm 0.3
PT Quadriceps/Body Mass				
60°s ⁻¹	3.4 \pm 0.4	3.3 \pm 0.5	3.4 \pm 0.5	3.4 \pm 0.5
240°s ⁻¹	2.1 \pm 0.2	2.2 \pm 0.4	2.1 \pm 0.3	2.2 \pm 0.3

Significant differences: * $p < 0.05$; ** $p < 0.01$ Nm: Newton meter PT: Peak Torque

Mean peak torque of the hamstring muscle of the left and right knees was significantly lower in soccer players compared with basketball players at 60°s⁻¹ ($p < 0.01$). Furthermore, mean peak torque was significantly lower in soccer players than basketball players at 240°s⁻¹ in the right knee ($p < 0.01$) and in the left knee ($p < 0.05$) (Table 2). Mean peak torque of the quadriceps muscle of the right knee was significantly higher in basketball players compared

to soccer players at 60°s^{-1} and 240°s^{-1} in the right ($p<0.01$) and the left knee ($p<0.05$) (Table 2).




FIGURE 1: HAMSTRING AND QUADRICEPS MUSCLE RATIO (%) AT PEAK TORQUE AT A VELOCITY OF 60°S^{-1}

There was only 1 significant between-group difference in peak torque H:Q muscle ratio. The left knee ratio at 60°s^{-1} (Figure 1) was significantly higher in the basketball players ($p<0.05$). There were no significant differences between soccer and basketball players in peak torque flexion/body mass ratio and in peak torque extension/body mass ratio (Table 2).

DISCUSSION

The study was conducted to determine and compare the maximal voluntary peak torques of the quadriceps and hamstring muscles, as well as on the reciprocal relationship between agonist and antagonist muscles of the right and left legs in professional soccer and basketball players. There were observed differences between soccer and basketball players. When these results were compared with those of other studies regarding peak torque of H and Q muscles, both similarities and differences in isokinetic peak torque H:Q muscle ratio and peak torque extension and flexion/body mass ratios were observed.

Regarding left and right knee extension and flexion, hamstring and quadriceps muscle peak torques were higher in basketball players than soccer players. Similar results have been found in previous studies (Zakas *et al.*, 1995; Metaxas *et al.*, 2009; Kazazovic *et al.*, 2010; Ozkan & Isler Kin, 2010; Alemdaroglu, 2012), that investigated maximal voluntary peak torques of the Q and H muscles and the torque ratio between these muscle groups in basketball and soccer players participating in teams of different divisions. Zakas *et al.* (1995) and Metaxas *et al.* (2009) found that peak torque expressed in absolute terms was significantly higher in basketball than soccer players at velocities of 60°s^{-1} and 180°s^{-1} .

Kazazovic *et al.* (2010) used isokinetic dynamometer measurements to evaluate maximum torque, total work and agonist–antagonist reciprocity of the knee joint in professional and amateur athletes at angular velocities of 60°s^{-1} and 180°s^{-1} . Their results revealed that professional athletes presented significantly higher values for total work and maximum torque of the knee flexors at an angular velocity of 60°s^{-1} . Moreover, peak torque H:Q muscle ratio was much higher in the basketball players' left leg than the soccer players' left leg at 60°s^{-1} (Dauty *et al.*, 2007). Metaxas *et al.* (2009) examined and compared cardio-respiratory performance and isokinetic muscle strength at angular velocities of 60°s^{-1} , 180°s^{-1} , and 300°s^{-1} between Greek soccer and basketball players of different divisions before starting the training season. Regarding peak torque, only basketball players in the IIIrd and IVth divisions showed significantly higher values at 60°s^{-1} in the hamstrings.

When the average peak torque H:Q muscle ratios were compared between basketball and soccer players, no significant differences were found between the groups except for the left leg at a velocity of 60°s^{-1} (Table 2). Peak torque H:Q muscle in the left limb at 60°s^{-1} was significantly greater in basketball than soccer players. Similar studies (Zakas *et al.*, 1995; Masuda *et al.*, 2003; Patricia & Vassilios, 2003; Ergun *et al.*, 2004; Magalhaes *et al.*, 2004; Egan *et al.*, 2006; Zakas, 2006; Tabakovic *et al.*, 2009; Brughelli *et al.*, 2010), yielded the same results. Brughelli *et al.* (2010) researched the effects of eccentric exercise on optimum length of the knee flexors and extensors during the preseason in professional soccer players. They reported that peak torque levels and H/Q ratios were not significantly altered throughout their study and that eccentric exercise could also increase the optimum lengths of both knee extensors and flexors during the preseason in professional soccer.

Ergun *et al.* (2004) conducted a cross-sectional analysis of sagittal knee laxity and isokinetic muscle strength in soccer players. Their results showed that the dominant extremity in soccer players had significantly higher knee flexor peak torque and H:Q muscle ratio at 180°s^{-1} and that those soccer players had significantly higher extensor and flexor peak torque values and H:Q muscle ratios than sedentary participants for both extremities.

However, Magalhaes *et al.* (2004) studied the concentric quadriceps and hamstring isokinetic strengths in volleyball and soccer players and reported that the H:Q muscle ratio was significantly lower in volleyball players at 90°s^{-1} and that no significant differences were found for H:Q muscle ratio in soccer players of different positional roles.

Additionally, Masuda *et al.* (2003) examined the relationships between muscle cross-sectional area (CSA) and muscular strength in terms of knee extension and flexion, hip extension and flexion and hip abduction and adduction among 14 well-trained university soccer players, who were divided into two groups based on ability (Group A: above-average ability; Group B: average ability). They observed no significant differences between the two groups in muscle CSA and isokinetic strength. Rahnama *et al.* (2003) studied how exercise that simulates the work rate of competitive soccer players affects the strength of the knee extensors and knee flexors and showed that significant changes were found ($p < 0.05$) for both legs in the hamstring to quadriceps muscle ratio.

No significant differences were found for average peak torque H/body mass ratio and Q/body mass ratio between the soccer and basketball players in this study. This indicates that soccer

and basketball players have the same relative strength. However, some researchers (Williams & Singh, 1997; Pincivero *et al.*, 2002; Buchanan & Vardaxis, 2003), found significant relationships between peak torque Q/body mass ratio and H/body mass ratio. Pincivero *et al.* (1997) studied the relationship between open and closed kinematic chain assessment of knee strength and functional performance. They noted that correlation coefficients were statistically greater for peak torque/body mass at a velocity of 180°s^{-1} .

Additionally, Williams and Singh (1997) studied dynamic trunk strength of Canadian football players, soccer players, and middle- to long-distance runners. According to their results, eccentric flexor peak torque relative to body mass was significantly greater in soccer players than runners and recreationally active participants. Buchanan and Vardaxis (2003; 2009) explored the differences in Q and H strength among the genders and different age groups and found that, with body mass-stature normalisation, most age and gender differences were small. Pincivero *et al.* (2002) assessed isokinetic torque, work and power among non-injured, ACL (anterior cruciate ligament)-deficient and ACL-reconstructed individuals. The H peak torque corrected for body mass was significantly higher in the non-involved than in the involved limb only at 60°s^{-1} .

CONCLUSION

This study revealed significant differences between professional basketball and soccer players for peak torques of H and Q muscles. However, when peak torque H:Q muscles ratio was compared between soccer and basketball players, there was only a significant difference for the left knee of basketball players at 60°s^{-1} . The difference was due to bilateral left- and right-leg strength differences in soccer players, which were not found in basketball players. Therefore, the isokinetic strength of the left knee might be lower than that of the right knee in soccer players. Strength training, especially for the left leg, may be necessary to reduce injury risk in soccer players. However, there were no significant differences in average peak torque Q/body mass and H/body mass ratios between basketball and soccer players.

Therefore, relative strength is comparable in soccer and basketball players, but absolute strength is greater in basketball than soccer players. According to the study data, body mass has a decisive effect on Q and H muscle peak torque values in soccer and basketball players and relationships also exist between mass and peak power.

Acknowledgement

The physical and physiological tests were carried out at the School of Physical Education and Sport at Balikesir University.

REFERENCES

- AAGAARD, P.; SIMONSEN, E.B.; TROLLE, M.; BANGSBO, J. & KLAUSEN, K. (1995). Isokinetic hamstring/quadriceps strength ratio: Influence from joint angular velocity, gravity correction and contraction mode. *Acta Physiologica Scandinavica*, 154: 421-427.

- ALEMDAROGLU, U. (2012). The relationship between muscle strength, anaerobic performance, agility, sprint ability and vertical jump performance in professional basketball players. *Journal of Human Kinetics*, 31: 149-158.
- BARNETT, L.; VAN BEURDEN, E.; MORGAN, P.; BROOKS, L. & BEARD, J. (2008). Does childhood motor skill proficiency predict adolescent fitness? *Medicine Science in Sports Exercise*, 40(12): 2137.
- BROCKETT, C.; MORGAN, D.L. & PROSKE, U. (1999). Using isokinetic dynamometry to indicate damage from eccentric exercise in human hamstring muscles. Paper presented at the 5th IOC World Congress on Sport Sciences, Sydney, Australia, 31 October - 5 November, Sydney, NSW, Australia, p. 31.
- BRUGHELLI, M.; MENDIGUCHIA, J.; NOSAKA, K.; IDOATE, F.; ARCOS, A.L. & CRONIN, J. (2010). Effects of eccentric exercise on optimum length of the knee flexors and extensors during the preseason in professional soccer players. *Physical Therapy in Sport*, 11(2): 50-55.
- BUCHANAN, P.A. & VARDAXIS, V.G. (2003). Sex-related and age-related differences in knee strength of basketball players ages 11-17 years. *Journal of Athletic Training*, 38: 231-237.
- BUCHANAN, P.A. & VARDAXIS, V.G. (2009). Lower-extremity strength profiles and gender-based classification of basketball players ages 9-22 years. *Journal of Strength and Conditioning Research*, 23: 406-419.
- CLANTON, T.O. & COUPE, K.J. (1998). Hamstring strain in athletes: Diagnosis and treatment. *Journal of the American Academy of Orthopaedic Surgeons*, 6: 237-248.
- COMETTI, G.; MAFFIULETI, N.A.; POUSSON, M.; CHATARD, J.C. & MAFFULLI, N. (2001). Isokinetic strength and anaerobic power of elite, sub-elite and amateur French soccer players. *International Journal of Sports Medicine*, 22: 45-51.
- CROISIER, J.L.; FORTHOMME, B.; NAMUROIS, M.H.; VANDERTHOMMEN, M. & CRIELAARD, J.M. (2002). Hamstring muscle strain recurrence and strength performance disorders. *American Journal of Sports Medicine*, 30(2): 199-203.
- CROISIER, J.L.; GANTEAUME, S.; BINET, J.; GENTY, M. & FERRET, J.M. (2008). Strength imbalances and prevention of hamstring injury in professional soccer players: A prospective study. *American Journal of Sports Medicine*, 36: 1469-1475.
- DAUTY, M.; DUPRE, M.; POTIRON-JOSSE, M. & DUBOIS, C. (2007). Identification of mechanical consequences of jumper's knee by isokinetic concentric torque measurement in elite basketball players. *Isokinetics and Exercise Science*, 15: 37-41.
- EGAN, A.D.; CRAMER, J.T.; MASSEY, L.L. & MAREK, S.M. (2006). Acute effects of static stretching on peak torque and mean power output in National Collegiate Athletic Association Division I women's basketball players. *Journal of Strength and Conditioning Research*, 20(4): 778-782.
- ERGUN, M.; ISLEGEN, C. & TASKIRAN, E. (2004). A cross-sectional analysis of sagittal knee laxity and isokinetic muscle strength in soccer players. *International Journal of Sports Medicine*, 25(8): 594-598.
- GRIFFIN, L.Y.; AGEL, J.; ALBOHM, M.J.; ARENDT, E.A.; DICK, R.W.; GARRETT, W.E.; GARRICK, J.G.; HEWETT, T.E.; HUSTON, L.; IRELAND, M.L.; JOHNSON, R.J.; KIBLER, W.B.; LEPHART, S.; LEWIS, J.L.; LINDENFELD, T.N.; MANDELBAUM, B.R.; MARCHAK, P.; TEITZ, C.C. & WOJTYNS, E.M. (2000). Noncontact anterior cruciate ligament injuries: Risk factors and prevention strategies. *Journal of the American Academy of Orthopaedic Surgeons*, 8: 141-150.
- HEWETT, T.E.; LINDENFELD, T.N.; RICCOBENE, J.V. & NOYES, F.R. (1999). The effect of neuromuscular training on the incidence of knee injury in female athletes: A prospective study. *American Journal of Sports Medicine*, 27: 699-706.

- KANNUS, P. (1988). Ratio of hamstrings to quadriceps femoris muscles' strength in the anterior cruciate ligament insufficient knee: Relationship to long-term recovery. *Journal of Physical Therapy*, 69: 961-965.
- KAZAZOVIC, E.; HADZIKADUNIC, A. & KOZIC, V. (2008). Effects of additional exercise program performed with Biodex apparatus at the maximal strength of the dynamic stabilization of knee muscles in active handball players. Paper presented at the 4th International Symposium on Youth Sport, 14 – 16 November, Ljubljana, Slovenia.
- KAZAZOVIC, E.; KOZIC, V.; SOLAKOVIC, E. & SEBIC-ZUHRIC, L. (2009). The effects of isokinetic exercise program on the knee flexing strength: Sport scientific practical aspects. *International Scientific Journal of Kinesiology*, 6(1): 25-30.
- KAZAZOVIC, E.; TABAKOVIC, M.; TALOVIC, M.; ALIC, H.; JELESKOVIC, E. & MRKOVIC, R. (2010). Evaluation of knee muscles isokinetic evaluation between professional and amateur athletes first year students of the faculty of sport and physical education. *Homo Sporticus*, 2: 32-35.
- LI, R.C.; MAFFULLI, N.; HSU, Y.C. & CHAN, K.M. (1996). Isokinetic strength of the quadriceps and hamstrings and functional ability of anterior cruciate deficient knees in recreational athletes. *British Journal of Sports Medicine*, 30: 161-164.
- MAGALHAES, J.; OLIVEIRA, J.; ASCENSAO, A. & SOARES, J. (2004). Concentric quadriceps and hamstrings isokinetic strength in volleyball and soccer players. *Journal of Sports Medicine and Physical Fitness*, 44: 119-125.
- MASUDA, K.; KIKUHARA, N.; TAKAHASHI, H. & YAMANAKA, K. (2003). The relationship between muscle cross-sectional area and strength in various isokinetic movements among soccer players. *Journal of Sports Science*, 21(10): 851-858.
- METAXAS, T.I.; KOUTLIANOS, N.; SENDELIDES, T. & MANDROUKAS, A. (2009). Preseason physiological profile of soccer and basketball players in different divisions. *Journal of Strength and Conditioning Research*, 23: 1704-1713.
- MOORE, J.R. & WADE, G. (1989). Prevention of anterior cruciate ligament injuries. *Journal of the National Strength and Conditioning Association*, 11: 35-40.
- OBERG, B.; MOLLER, M.; GILLQUIST, J. & EKSTRAND, J. (1986). Isokinetic torque levels for knee extensors and knee flexors in soccer players. *International Journal of Sports Medicine*, 7(1): 50-53.
- ORCHARD, J.; MARSDEN, J.; LORD, S. & GARLICK, D. (1997). Preseason hamstring muscle weakness associated with hamstring muscle injury in Australian footballers. *American Journal of Sports Medicine*, 25: 81-85.
- OZKAN, A. & ISLER KIN, A. (2010). The association among leg volume, leg mass and H:Q ratio with anaerobic performance and isokinetic knee strength in athletes. *Hacettepe University Journal of Sport Science*, 21(3): 90-102.
- PANNI, A.; BIEDERT, R.M.; MAFFULLI, N.; TARTARONE, M. & ROMANINI, E. (2002). Overuse injury to the extensor mechanism in athletes. *Clinical Sports Medicine*, 21: 483-498.
- PATRICIA, A.B. & VASSILIOS, G.V. (2003). Sex-related and age-related differences in knee strength of basketball players ages 11-17 years. *Journal of Athletic Training*, 38(3): 231-237.
- PERRIN, D.H.; ROBERTSON, R.J. & RAY, R.L. (1987). Bilateral isokinetic peak torque, torque acceleration energy, power, and work relationships in athletes and nonetheless. *Journal of Orthopedic and Sports Physical Therapy*, 9: 184-189.
- PETERSEN, J. & HOLMICH, P. (2005). Evidence based prevention of hamstring injuries in sport. *British Journal of Sports Medicine*, 39: 319-323.

- PETTITT, R.W. & BRYSON, E.R. (2002). Training for women's basketball: A biomechanical emphasis for preventing anterior cruciate ligament injury. *Strength and Conditioning Journal*, 24: 20-29.
- PINCIVERO, D.M.; LEPHART, S.M. & KARUNAKARA, R.G. (1997). Relation between open and closed kinematic chain assessment of knee strength and functional performance. *Clinical Journal of Sport Medicine*, 7(1): 11-18
- PINCIVERO, D.M.; HELLER, B.M. & HOU, S.I. (2002). The effects of ACL injury on quadriceps and hamstring torque, work and power. *Sports Sciences*, 20(9): 689-696.
- RAHNAMA, N.; REILLY, T.; LEES, A. & GRAHAM-SMITH, P. (2003). Muscle fatigue induced by exercise simulating the work rate of competitive soccer. *Journal of Sports Science*, 21(11): 933-942.
- REILLY, T. & THOMAS, V. (1976). A motion analysis of work-rate in different positional roles in professional football match-play. *Journal of Human Movement Studies*, 2: 87-89.
- RICHARDS, D.P.; AJEMIAN, S.V.; WILEY, P. & ZERNICKE, R.F. (1996). Knee joint dynamics predict patellar tendinitis in elite volleyball players. *American Journal of Sports Medicine*, 24: 676-683.
- SIQUEIRA, C.M.; PELLEGRINI, F.R.; FONTANA, M.F. & GREVE, J.M. (2002). Isokinetic dynamometry of knee flexors and extensors: Comparative study among non-athletes, jumper athletes and runner athletes. *Revista do Hospital das Clinicas Faculty Medicine (Sao Paulo)*, 57: 19-24.
- TABAKOVIC, M.; KAZAZOVIC, E.; TALOVIC, M. & TURKOVIC, S. (2009). Bilateral and reciprocal relation between extensor and flexor knee strength in football and basketball players. *Homo Sporticus*, 1: 33-36.
- THISSEN-MILDER, M. & MAYHEW, J.L. (1990). Selection and classification in high school volleyball players from performance tests. *Journal of Sports Medicine and Physical Fitness*, 31: 380-384.
- WILLIAMS, C.A. & SINGH, M. (1997). Dynamic trunk strength of Canadian football players, soccer players, and middle to long distance runners. *Journal of Orthopaedic and Sports Physical Therapy*, 25(4): 271-276.
- ZAKAS, A. (2006). Bilateral isokinetic peak torque of quadriceps and hamstring muscles in professional soccer players with dominance on one or both two sides. *Journal of Sports Medicine and Physical Fitness*, 46 (1):28-35.
- ZAKAS, A.; MANDROUKAS, K.; VAMVAKOUDIS, E.; CHRISTOULAS, K. & AGGELOPOULOU, N. (1995). Peak torque of quadriceps and hamstring muscles in basketball and soccer players of different divisions. *Journal of Sports Medicine and Physical Fitness*, 35: 199-205.

Assist. Prof Dr Ibrahim ERDEMİR: School of Physical Education and Sports, Dinkçiler Mahallesi Soma Caddesi 10100, Balıkesir University, Balıkesir, Turkey. Tel.: +90 266 238 18 38, Fax.: +90 266 239 02 85, Cell: +90 532 227 19 30, E-mail: iboerdemir@gmail.com

(Subject Editor: Prof Ernst Krüger)